

Set rules:

Let t, t' stand for any closed term, y for any variable:

$$\frac{\frac{\neg St}{\forall y y \notin t} \quad \frac{t \in t'}{St'}}{S} \quad S$$

7. 1 EXTENSIONALITY

$$\forall x \forall y ((Sx \ \& \ Sy) \rightarrow (\forall z (z \in x \leftrightarrow z \in y) \rightarrow x = y))$$

Derivable rule.

Let t and t' stand for any closed term, v for any variable:

$$\frac{St \quad St'}{\forall v (v \in t \leftrightarrow v \in t')}$$

$$t = t'$$

Ext.

⊆ Exploitation

$$\frac{\underline{t \subseteq t'}}{St \ \& \ St' \ \& \ \forall v(v \in t \rightarrow v \in t')} \quad \underline{\subseteq E}$$

$$\frac{\underline{\neg St}}{\neg t \subseteq t'} \quad \underline{\subseteq E}$$

$$\frac{\underline{\neg St'}}{\neg t \subseteq t'} \quad \underline{\subseteq E}$$

New formation rule:

If v is a variable, t is a closed term, c is a constant, \mathcal{A} is a formula containing c but not containing v , then:

$\{v \in t: \mathcal{A} [v/c] \}$ is a closed term.

Such a term is called an abstraction term. ' v ' is the variable of abstraction, ' t ' is the restrictor term, and $\mathcal{A} [v/c]$ is the abstraction formula.

Abstraction axiom schema:

$$\forall x (S\{z \in x : \mathcal{A}(z)\} \ \& \ \forall y(y \in \{z \in x : \mathcal{A}(z)\} \leftrightarrow (y \in x \ \& \ \mathcal{A}(y))))$$

Here $\mathcal{A}(z) = \mathcal{A}[z/c]$ for some \mathcal{A} containing c and not z . Likewise, $\mathcal{A}(y) = \mathcal{A} [y/c]$.

Theorem 7.4: Any set x is the set of its self-identical members, $\{y \in x: y = y\}$

1. Show $\forall x(Sx \rightarrow (x = \{y \in x: y = y\}))$
2. Show $Sa \rightarrow (a = \{y \in a: y = y\})$
3. Sa
4. $S\{y \in a: y = y\} \&$
 $\forall z(z \in \{y \in a: y = y\} \leftrightarrow$
 $(z \in a \& z = z))$ Abst.
5. $\forall z(z \in \{y \in a: y = y\} \leftrightarrow$
 $(z \in a \& z = z))$ &E, 4
6. Show $\forall z(z \in a \leftrightarrow z \in \{y \in a: y = y\})$
7. Show $c \in a \leftrightarrow c \in \{y \in a: y = y\}$
8. $c \in \{y \in a: y = y\} \leftrightarrow (c \in a \& c = c)$
 $\forall E, 5$
9. Show $c \in a \rightarrow c \in \{y \in a: y = y\}$
10. $c \in a$
11. $c = c$ =I
12. $c \in a \& c = c$ 10,11,&I
13. $c \in \{y \in a: y = y\}$ $\leftrightarrow E, 8, 12$
14. Show $c \in \{y \in a: y = y\} \rightarrow c \in a$
15. $c \in \{y \in a: y = y\}$

16. $c \in a \ \& \ c = c$ $\leftrightarrow E, 8, 15$
 17. $\frac{\frac{\frac{}{c \in a}}{}{}{}}{}{}$ $16, \ \&E$
 18. $S\{y \in a: y = y\}$ $4, \ \&E$
 19. $a = \{y \in a: y = y\}$ $3, 18, 6, \text{Ext}$

Theorem 7.3: $\forall x S\{z \in x: A(z)\}$

Let v be any variable, t and t' any closed terms.

Let $\mathcal{A}(v) = \mathcal{A}[v/c]$, where \mathcal{A} is some formula containing c and not v .

Let $\mathcal{A}(t) = \mathcal{A}[t/c]$.

Naive Abstraction:

$$\frac{\frac{t \in \{v \in t': A(v)\}}{}{}}{t \in t' \ \& \ A(t)} \quad \text{Abstr.}$$

1. Show $\forall x \exists y y \notin x$
2. Show $\exists y y \notin a$
3. Show $\overline{\{z \in a: z \notin z\}} \notin a$
4. $\{z \in a: z \notin z\} \in a$
5. $\{z \in a: z \notin z\} = b$ =I, \exists I, \exists E
6. Show $b \in b$
7. $b \notin b$
8. $b \in a$ =E, 4, 5
9. $b \in a \ \& \ b \notin b$ &I, 7, 8
10. $b \in \{z \in a: z \notin z\}$ Abst., 9
11. $b \in b$ =E, 4, 10
12. $b \in \{z \in a: z \notin z\}$ =E, 5, 6
13. $b \in a \ \& \ b \notin b$ Abst., 12
14. $b \notin b$ &E, 13
15. $\overline{\exists y y \notin a}$ \exists I, 3

New formation rules:

If t and t' are closed terms, then $\{t, t'\}$ is a closed term.

If t_1, \dots, t_i are closed terms, then $\{t_1, \dots, t_i\}$ is a closed term.

Unit Set

$$\underline{\underline{x \in \{y\}}}$$

$$x = y$$

Unit

Pair Set

$$\underline{\underline{u \in \{y, z\}}}$$

$$u = y \vee u = z$$

Pair set

Enumeration

$$\underline{\underline{x \in \{y_1, \dots, y_i\}}}$$

$$x = y_1 \vee x = y_2 \vee \dots \vee x = y_i$$

Enum

Everything belongs to some set.

1. Show $\forall x \exists y (Sx \& x \in y)$

2. Show $\exists y (Sy \& a \in y)$

3. $S\{a\}$

Thm 6

4. $a \in \{a\}$

Unit E*

5. $Sa \& a \in \{a\}$

6. $\exists y (Sy \& a \in y)$

$\exists I, 5$

Theorem 7.10. There is a set without members. $\exists x (Sx \ \& \ \forall y \ y \notin x)$

- | | | |
|----|--|--------------|
| 1. | Show $\exists x(Sx \ \& \ \forall y (y \notin x))$ | |
| 2. | $S\{x: x \neq x\}$ | Th 7.3 |
| 3. | <u>Show</u> $\forall y \ y \notin \{x: x \neq x\}$ | |
| 4. | <u>Show</u> $c \notin \{x: x \neq x\}$ | |
| 5. | $c \in \{x: x \neq x\}$ | AIP |
| 6. | $c \neq c$ | Abst, 5 |
| 7. | $c = c$ | =I |
| 8. | $\exists x (Sx \ \& \ \forall y \ y \notin x)$ | &I, 2, 3, EI |

Definition: $\emptyset = \{x: x \neq x\}$

Derivable inference rule:

$$\frac{\emptyset}{t \notin \emptyset} \quad \emptyset$$

Theorem 7.11 $S\emptyset \ \& \ \forall y \ y \notin \emptyset$

Theorem 7.12. \emptyset is the only empty set.

$$\forall x((Sx \ \& \ \forall y \ y \notin x) \rightarrow x = \emptyset)$$

1. Show $\forall x((Sx \ \& \ \forall y \ y \notin x) \rightarrow x = \emptyset)$
2. Show $(Sa \ \& \ \forall y \ y \notin a) \rightarrow a = \emptyset$
3. $Sa \ \& \ \forall y \ y \notin a$ ACP
4. Show $\forall z(z \in a \leftrightarrow z \in \emptyset)$
5. Show $b \in a \leftrightarrow b \in \emptyset$
6. Show $b \in a \rightarrow b \in \emptyset$
7. $b \in a$ ACP
8. $b \notin a$ $\forall E3, \ \&E \ 3$
9. $b \in \emptyset$!, 7, 8
10. Show $b \in \emptyset \rightarrow b \in a$
11. $b \in \emptyset$ ACP
12. $b \notin \emptyset$ \emptyset
13. $b \in a$!, 11, 12
14. $b \in a \leftrightarrow b \in \emptyset$ $\leftrightarrow I, \ 6, 10$
15. $a = \emptyset$ Ext, 4, 3, Thm 7.11

Corollary 7.1

$$\forall x ((Sx \ \& \ x \neq \emptyset) \rightarrow \exists y \ y \in x)$$